

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
24 June 2004 (24.06.2004)

PCT

(10) International Publication Number  
**WO 2004/052881 A2**

(51) International Patent Classification<sup>7</sup>: **C07D 401/12**

(81) Designated States (*national*): AE, AI, AU, BA, BR, CA, CN, CO, DZ, EC, EG, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, US, VN, YU, ZA, ZW.

(21) International Application Number:  
PCT/EP2003/013604

(22) International Filing Date: 3 December 2003 (03.12.2003)

(84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
02027274.6 6 December 2002 (06.12.2002) EP  
103 40 254.3 29 August 2003 (29.08.2003) DE

**Declarations under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AL, AU, BA, BR, CA, CN, CO, DZ, EC, EG, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, VN, YU, ZA, ZW; Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR)

— of inventorship (Rule 4.17(iv)) for US only

(71) Applicant (*for all designated States except US*): ALTANA PHARMA AG [DE/DE]; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): KOHL, Bernhard [DE/DE]; Zum Brühl 9, 78465 Konstanz (DE). MÜLLER, Bernd [DE/DE]; Bücklestr. 84a, 78467 Konstanz (DE). WEINGART, Ralf Steffen [DE/DE]; Thingoltstr. 34, 78465 Konstanz (DE).

**Published:**

— without international search report and to be republished upon receipt of that report

(74) Agent: WOLF, Ulrich; Altana Pharma AG, Byk-Gulden-Str. 2, 78467 Konstanz (DE).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PROCESS FOR PREPARING (S)-PANTOPRAZOLE

(57) Abstract: The invention relates to a novel process for preparing (S)-pantoprazole using a chiral zirconium complex or a chiral hafnium complex.

WO 2004/052881 A2

## Process for preparing (S)-pantoprazole

### Subject-matter of the invention

The present invention relates to a novel process for preparing the active compound (S)-pantoprazole which can be used for preparing medicaments in the pharmaceutical industry.

### Technical background

Pyridin-2-ylmethylsulphonyl-1H-benzimidazoles and compounds of a closely related structure, as known, for example, from EP-A-0005129, EP-A-0166287, EP-A-0174726 and EP-A-0268956, are, owing to their  $H^+/K^+$ -ATPase-inhibitory action, of considerable importance in the therapy of diseases associated with an increased secretion of gastric acid.

Examples of active compounds from this class of compounds which are commercially available or in clinical development are 5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: omeprazole), (S)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: esomeprazole), 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: pantoprazole), 2-[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: lansoprazole), 2-[[4-(3-methoxypropoxy)-3-methylpyridin-2-yl)methylsulphonyl]-1H-benzimidazole (INN: rabeprazole) and 5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridylmethylsulphonyl)-1H-imidazo(4,5-b)pyridine (INN: tenatoprazole).

The abovementioned sulphonyl derivatives which, owing to their mechanism of action, are also referred to as proton pump inhibitors or abbreviated PPI are chiral compounds. The process usually used for preparing the PPI is the oxidation of the corresponding sulphides. This oxidation gives – unless particular measures are taken – a racemic mixture comprising about the same proportions of the two enantiomers (stereoisomers), i.e. the (+)- and (-)-form or the (R)- and (S)-form of the PPI.

Since enantiomers are thermally relatively stable, i.e. they do not racemize on storage – in particular in solid form – there has in the past been no lack of efforts to separate PPI enantiomer mixtures or to prepare the PPI enantiomers in more or less pure form.

### Prior art

The international patent application WO91/12221 describes a process for separating enantiomers using a cellulase enzyme. One of the active compounds mentioned as being separable into the enantiomers with the aid of this process is omeprazole.

The international patent application WO92/08716 describes, for the first time, a chemical process which allows the separation of pyridin-2-ylmethylsulphonyl-1H-benzimidazoles into their optical isomers. Compounds mentioned as having been prepared in an exemplary manner are, inter alia, the

compounds (+)- and (-)-5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole [= (+)- and (-)-pantoprazole]. The international patent application WO92/08716 refers to the fact that the optical isomers of the pyridin-2-yl-methylsulphonyl-1H-benzimidazoles, i.e. the (+)- and (-)-enantiomers or the (R)- and (S)-enantiomers, are used as active compounds in medicaments for the treatment of gastrointestinal disorders. With respect to the mode of application and the dosage of the active compounds, reference is made inter alia to the European patent 166 287.

The international patent application WO94/27988 describes the separation of racemic omeprazole into the enantiomers, using chiral auxiliaries.

The international patent application WO96/02535 (= USP 5,948,789) describes a process for the enantioselective synthesis of PPI using chiral titanium complexes. What is described is, inter alia, the synthesis of (+)- and (-)- [or, expressed in a different way, (R)- and (S)]-pantoprazole, the chiral auxiliary used for the synthesis of (+)-pantoprazole being diethyl (+)-tartrate and the chiral auxiliary used for the preparation of (-)-pantoprazole being diethyl (-)-tartrate.

The international patent applications WO96/17076 and WO96/17077 describe the enantioselective biooxidation or bioreduction with the use of certain microorganisms for the preparation of enantiomerically pure or enantiomerically enriched PPI.

The international patent application WO97/02261 describes the enrichment of PPI enantiomers by selective precipitation.

The international patent applications WO94/24867 and WO94/25028 claim the use of the compounds (-)- and (+)-pantoprazole for treating stomach disorders in humans. Each of the stereoisomers is said to have medical advantages compared to the respective other stereoisomers.

The enantioselective sulfoxidation for preparing esomeprazole ((S)-omeprazole) on a large scale using a chiral titanium complex is described in Tetrahedron, Asymmetry, (2000), 11, 3819-3825.

The enantioselective sulfoxidation of aryl alkyl sulphides and dialkyl sulphides in the presence of a zirconium catalyst having a polydentate ligand is described in J. Org. Chem., (1999), 64(4), 1327.

### **Description of the invention**

The invention provides a process for preparing (-)- or (S)-pantoprazole. The process is characterized in that the oxidation of the corresponding sulphide is carried out in the presence of a chiral zirconium complex or a chiral hafnium complex, the chiral auxiliary used being a (+)-L-tartaric acid derivative.

The fact that, when a chiral zirconium complex or a chiral hafnium complex is used for preparing (-)- or (S)-pantoprazole, it is possible to use, as chiral auxiliary, preferably a (+)-L-tartaric acid derivative instead of a (-)-D-tartaric acid derivative is surprising and particularly advantageous, since (+)-L-tartaric acid derivatives - with respect to naturally considerably more frequently occurring (+)-L-tartaric acid - are considerably less expensive and as a consequence highly suitable in particular for a preparation on an industrial scale.

The oxidation is advantageously carried out in an organic solvent, such as, for example, ethyl acetate, toluene, dichloromethane, dioxane or, preferably, methyl isobutyl ketone, where it is not necessary for the solvents mentioned to be completely anhydrous or where anhydrous solvents are in each case optionally admixed with a defined proportion of water, for example up to a maximum of 0.5 equivalent. For reactions with less than 0.5 equivalent of zirconium or hafnium complex, it is preferred to use an anhydrous solvent. The solvents employed may be used in the commercially available quality.

A solvent essentially comprises a specific solvent if it contains at least 50%, preferably at least 90%, in particular at least 95%, of the said specific solvent. An anhydrous solvent is essentially free of water, having a water content of less than 5%, preferably less than 1%, in particular less than 0.3%.

Suitable oxidizing agents are all anhydrous oxidizing agents customarily used for the synthesis of PPI, where particular mention may be made of hydroperoxides, such as, for example, tert-butyl hydroperoxide or, in particular, cumene hydroperoxide. In general, 0.90 to 1.3 oxidation equivalents, preferably 0.95-1.05 equivalents, of the oxidizing agent are used.

Suitable zirconium complexes are, for example, zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide and, in particular, zirconium(IV) n-propoxide (preferably as a solution in n-propanol) or zirconium(IV) isopropoxide (preferably in the form of the zirconium(IV) isopropoxide/isopropanol complex). Suitable hafnium complexes are, for example, hafnium(IV) acetylacetonate, hafnium(IV) butoxide, hafnium(IV) n-propoxide, hafnium(IV) isopropoxide (preferably in the form of the hafnium(IV) isopropoxide/isopropanol complex), hafnium(IV) ethoxide and in particular hafnium(IV) tert-butoxide. Preference is given to using a zirconium complex.

In general, 0.01-2 equivalents, preferably 0.05-0.9 equivalent, of the zirconium complex or of the hafnium complex are used.

Suitable (+)-L-tartaric acid derivatives are, for example, tartaric acid amides, such as (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide) or (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), or dialkyl tartrates,

such as dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate and diethyl (+)-L-tartrate. In general, 0.02-4 equivalents, preferably 0.1-2 equivalents, of the (+)-L-tartaric acid derivative are employed.

Particularly preferred tartaric acid derivatives are (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-morpholinamide).

The oxidation is preferably carried out at temperatures between -20 and 50°C, in particular at room temperature, and optionally in the presence of a base, suitable bases being, in particular, organic bases, preferably a tertiary amine, such as triethylamine or N-ethyldiisopropylamine.

If the process is carried out in a suitable manner, (-)- or (S)-pantoprazole is obtained in an optical purity of >95%. By further steps, such as, for example, pH-controlled reprecipitation and/or recrystallization in a suitable solvent, such as, for example, isopropanol, it is possible to further increase the optical purity considerably. Reprecipitation is carried out via intermediate preparation of suitable salts, such as, for example, via the sodium salt (for other possible salts, see, for example, EP-A-166287).

The invention is illustrated in more detail by the examples below, but not limited in any way. The abbreviation h stands for hour(s).

### Examples

**1. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulfinyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with diethyl (+)-L-tartrate and zirconium(IV) isopropoxide/isopropanol**

A) At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole together with 17.9 g of diethyl (+)-tartrate, 13.4 g of zirconium(IV) isopropoxide/isopropanol and 0.1 ml of water are suspended in 100 ml of methyl isobutyl ketone. The mixture is heated at 40°C for one hour, resulting in the formation of an almost clear solution. After cooling to room temperature, 4.1 ml of N-ethyldiisopropylamine are added. 11 ml of cumene hydroperoxide are then slowly metered in. Stirring at room temperature is continued until the oxidation process has ended (monitored by TLC). The clear solution is quenched with 0.9 g of sodium thiosulphate in 54 ml of water and 30.3 g of 40% (w/w) of NaOH and stirred for another 14 h. After addition of 25 g of sodium chloride, the phases are separated. The aqueous phase is extracted with 50 ml of methyl isobutyl ketone. The combined organic phases are washed together using 25 ml of saturated sodium chloride solution. 150 ml of water are added to the methyl isobutyl ketone solution, and the pH is adjusted to 13 using 10% (w/w) NaOH. The phases are separated and the methyl isobutyl ketone phase is extracted once more with 50 ml of water at pH 13. The aqueous phases are combined and, at 40°C and under reduced pressure, subjected to incipient distillation. At 40-50°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH 9. Under pH control, stirring is continued for another 12 h. The beige crystals are filtered off and washed with 50 ml of water. This gives the title compound in an optical purity of >90%.

To increase the purity, (-)-pantoprazole is dissolved in water/NaOH and again precipitated by addition of acetic acid to pH 9. Drying gives a beige powder of melting point 145°C (decomposition) and an optical purity of >95%. If this powder is recrystallized from 2-PrOH, a clear crystal of melting point 147-149°C (decomposition) with an optical rotation of  $\alpha_D^{20} = -140$  (c=0.5, MeOH) is obtained.

B) Alternatively, the reaction described in Example 1A can be carried out in 100 ml of toluene instead of methyl isobutyl ketone. If the reaction is carried out in toluene, the zirconium salts have to be filtered off after quenching and the reaction product ((S)-pantoprazole as sodium salt) is directly extracted into the aqueous phase. From the aqueous phase, it can then be precipitated under controlled pH as (S)-pantoprazole. This gives beige crystals of an optical purity of > 95%.

**2. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-dimethylamide) and zirconium(IV) isopropoxide/isopropanol**

At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole are suspended in 100 ml of methyl isobutyl ketone together with 18.0 g of (+)-L-tartaric acid bis-(N,N-dimethylamide) and 13.4 g of zirconium(IV) isopropoxide-isopropanol. The mixture is heated at 40°C for one hour, resulting in the formation of a solution which is almost clear. After cooling to room temperature, 4.1 ml of N-ethyl-diisopropylamine are added. 11 ml of cumene hydroperoxide are then slowly metered in. The mixture is stirred at room temperature until the oxidation has ended (5-10 hours, monitored by TLC). The clear solution is diluted with 100 ml of methyl isobutyl ketone and quenched with 1.8 g of sodium thiosulphate in 140 ml of water and stirred for a further 14 hours. After phase separation, 55 ml of saturated sodium bicarbonate solution and 55 ml of methyl isobutyl ketone are added to the aqueous phase, and the phases are separated. Another 55 ml of saturated sodium bicarbonate solution and 55 ml of methyl isobutyl ketone are added to the aqueous phase, and the phases are separated. The combined methyl isobutyl ketone phases are then washed twice with 55 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to pH = 13 using a 40% by weight strength aqueous solution of sodium hydroxide. After phase separation, the methyl isobutyl ketone phase is extracted with another 50 ml of water at pH = 13. The aqueous phases are combined and, at 40°C, subjected to incipient distillation under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. The mixture is stirred for another 12 hours during which the pH is monitored. The beige crystals are filtered off and washed with 50 ml of water. The title compound is obtained in a yield of about 15 g (73% of theory) and an optical purity of >95%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and re-precipitated with acetic acid (10%) at pH = 9.0.

**3. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide) and zirconium(IV) isopropoxide/isopropanol**

At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole are suspended in 100 ml of methyl isobutyl ketone together with 22.6 g of (2R,3R)-(+)-L-tartaric acid bis-(N-pyrrolidinamide) and 13.4 g of zirconium(IV) isopropoxide-isopropanol. The mixture is heated at 40°C for one hour, resulting in the formation of a solution which is almost clear. After cooling to room temperature, 4.1 ml of N-ethyl-diisopropylamine are added. 11 ml of cumene hydroperoxide are then slowly metered in. The mixture is stirred at room temperature until the oxidation has ended (5-10 hours, monitored by TLC). The clear solution is diluted with 100 ml of methyl isobutyl ketone and quenched with 1.8 g of sodium thiosulphate in 140 ml of saturated sodium bicarbonate solution and stirred for a further 14 hours. After phase separation, the mixture is washed twice with 55

ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to pH = 13 using a 40% by weight strength aqueous solution of sodium hydroxide. After phase separation, the methyl isobutyl ketone phase is extracted with another 50 ml of water at pH = 13. The aqueous phases are combined and, at 40°C, subjected to incipient distillation under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. The mixture is stirred for another 12 hours during which the pH is monitored. The beige crystals are filtered off and washed with 50 ml of water. The title compound is obtained in a yield of about 17 g (80% of theory) and an optical purity of >98%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and re-precipitated with acetic acid (10%) at pH = 9.0.

**4. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole  
[= (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide)  
and zirconium(IV) n-propoxide**

At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole are suspended in 100 ml of methyl isobutyl ketone together with 22.6 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and 16.5 g of zirconium(IV) n-propoxide (70% in propanol). The mixture is heated at 40°C for one hour, resulting in the formation of a solution which is almost clear. After cooling to room temperature, 4.1 ml of N-ethyldiisopropylamine are added. 10 ml of cumene hydroperoxide are then slowly metered in. The mixture is stirred at room temperature until the oxidation has ended (5-24 hours, monitored by TLC). The clear solution is diluted with 100 ml of methyl isobutyl ketone and quenched with 1.8 g of sodium thiosulphate in 140 ml of saturated sodium bicarbonate solution and stirred for a further 14 hours. After phase separation, the mixture is washed twice with 55 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to pH = 13 using a 40% by weight strength aqueous solution of sodium hydroxide. After phase separation, the methyl isobutyl ketone phase is extracted with another 50 ml of water at pH = 13. The aqueous phases are combined and, at 40°C, subjected to incipient distillation under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. The mixture is stirred for another 12 hours during which the pH is monitored. The beige crystals are filtered off and washed with 50 ml of water. The title compound is obtained in a yield of about 16 g (75% of theory) and an optical purity of >98%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and re-precipitated with acetic acid (10%) at pH = 9.0.



5. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide) and zirconium(IV) n-propoxide

Analogously to Example 4, reaction of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole under otherwise identical conditions, but without addition of N-ethyl-diisopropylamine, gives the title compound in a yield of 65% of theory and an optical purity of >98%.

6. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and zirconium(IV) n-propoxide

Analogously to Example 4, reaction of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole under otherwise identical conditions, but with 0.1 equivalent of zirconium n-propoxide, 0.25 equivalent of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and 0.07 equivalents of Hünig base gives, after an oxidation time of 48-72 h, the title compound in a yield of 80% of theory and an optical purity of >98%.

7. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and zirconium(IV) n-propoxide

At room temperature, 50.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 5.2 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) (0.15 eq.) are suspended in 360 ml of methyl isobutyl ketone (MIBK). The suspension is heated at 40-45°C and 60 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 3.2 g of zirconium(IV) n-propoxide (70% in propanol, 0.05 eq.) are added, and the mixture is stirred for 1 hour. After cooling to 30°C, 0.9 ml of N-ethyl-diisopropylamine are added. 27.1 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (20 hours, monitored by TLC or HPLC). The suspension is diluted with 60 ml of 2-propanol and quenched with 1.69 g of sodium thiosulphate in 100 ml of saturated sodium bicarbonate solution and stirred for at least 2 hours. After phase separation, the mixture is washed twice with 50 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to 12.5-13 using 10 ml of aqueous sodium hydroxide solution (40% (w/w)). After phase separation, the methyl isobutyl ketone phase is extracted 2 more times with 100 ml of water and 2 ml of aqueous sodium hydroxide solution (40% (w/w)) at pH = 12.5-13. The combined aqueous phases are reextracted twice with 50 ml of methyl isobutyl ketone and subjected to incipient distillation at 40°C under reduced pressure. At 40-45°C, (-)-pantoprazole is

precipitated by addition of 10% strength acetic acid to pH = 9.0. Under pH control, stirring is continued for another 12 hours. The beige crystals are filtered off and washed twice with 50 ml of water.

This gives the title compound in a yield of 82% of theory in a chemical purity of 95% and an optical purity of > 95%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%). This gives the title compound in a yield of 75% of theory in a chemical purity of > 97% and an optical purity of > 98%.

**8. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and zirconium(IV) isopropoxide/isopropanol**

At room temperature, 10.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 1.05 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) (0.15 eq.) are suspended in 72 ml of methyl isobutyl ketone. The suspension is heated at 40–45°C and 12 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 0.53 g of zirconium(IV) isopropoxide/isopropanol (0.05 eq.) is added and the mixture is stirred for 1 hour. After cooling to 30°C, 0.16 ml of N-ethyl-diisopropylamine is added. 5.5 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (20 hours, monitored by TLC or HPLC). HPLC of the reaction shows 82% of title compound in an optical purity of > 95%.

**9. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and zirconium(IV) n-propoxide**

At room temperature, 50.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 13.9 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) (0.40 eq.) are suspended in 360 ml of methyl isobutyl ketone. The suspension is heated at 40–45°C and 60 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 6.4 g of zirconium(IV) n-propoxide (70% in propanol, 0.10 eq.) are added, and the mixture is stirred for 1 hour. After cooling to 30°C, 1.8 ml of N-ethyl-diisopropylamine are added. 27.1 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (20 hours, monitored by TLC or HPLC: chemical purity: 90% of pantoprazole sulphoxide). The suspension is diluted with 120 ml of 2-propanol and quenched with 1.69 g of sodium thiosulphate in 100 ml of saturated sodium bicarbonate solution and stirred for at least 2 hours. After phase separation, the mixture is washed twice with 50 ml of saturated sodium bicarbonate solution. 350 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to 12.5–13 using 10 ml of aqueous sodium hydroxide solution (40% (w/w)). After phase separation, the methyl isobutyl

ketone phase is extracted 2 more times with 100 ml of water and 2 ml of aqueous sodium hydroxide solution (40% (w/w)) at pH = 12.5-13. The combined aqueous phases are reextracted twice with 50 ml of methyl isobutyl ketone and subjected to incipient distillation at 40°C under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. Under pH control, stirring is continued for another 12 hours. The beige crystals are filtered off and washed twice with in each case 50 ml of water.

This gives the title compound in a yield of 85% of theory in a chemical purity of 95% and an optical purity of > 95%. To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%). This gives the title compound in a yield of 75-80% of theory in a chemical purity of > 98% and an optical purity of > 99%.

**10. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide) and hafnium(IV) tert-butoxide**

At room temperature, 3.67 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole, 4.10 g of (+)-L-tartaric acid bis-(N,N-pyrrolidinamide) and 2.60 ml of hafnium(IV) tert-butoxide are suspended in 18.5 ml of methyl isobutyl ketone. The mixture is heated at 40°C for 1 hour, during which an almost clear solution is formed. After cooling to room temperature, 0.74 ml of N-ethyl-diisopropylamine is added. 2.2 ml of cumene hydroperoxide are then slowly metered in. Stirring is continued at room temperature until the oxidation process has ended (48 hours, monitored by TLC). The clear solution is diluted with 20 ml of methyl isobutyl ketone and quenched with 0.3 g of sodium thiosulphate in 25 ml of saturated sodium bicarbonate solution and stirred for a further 14 hours. After phase separation, the methyl isobutyl ketone phase is washed two more times with 10 ml of saturated sodium bicarbonate solution. 30 ml of water are added to the methyl isobutyl ketone, phase and the pH is adjusted to 13 using 40% strength (w/w) aqueous sodium hydroxide solution. After phase separation, the methyl isobutyl ketone phase is once more extracted with 10 ml of water at pH = 13. The aqueous phases are combined and, at 40°C and under reduced pressure, subjected to incipient distillation. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH 9.0. Under pH control, stirring is continued for another 12 hours. The beige crystals are filtered off and washed with 10 ml of water. This gives the title compound in a yield of 2.5 g (65% of theory) in an optical purity of > 95%. To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%).

**Patent claims**

1. Process for preparing (S)-pantoprazole in enantiomerically pure or enantiomerically enriched form by oxidation of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole, characterized in that the oxidation is carried out in the presence of a chiral zirconium complex or a chiral hafnium complex.
2. Process for preparing (S)-pantoprazole in enantiomerically pure or enantiomerically enriched form by oxidation of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole, characterized in that the oxidation is carried out in the presence of a chiral zirconium complex.
3. Process according to Claim 1, characterized in that (S)-pantoprazole is obtained in an optical purity of > 90%.
4. Process according to Claim 1, characterized in that the oxidation is carried out using cumene hydroperoxide.
5. Process according to Claim 1, characterized in that zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex or hafnium(IV) acetylacetonate, hafnium(IV) butoxide, hafnium(IV) tert-butoxide, hafnium(IV) ethoxide, hafnium(IV) n-propoxide, hafnium(IV) isopropoxide or hafnium(IV) isopropoxide/isopropanol complex is used.
6. Process according to Claim 2, characterized in that zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex is used.
7. Process according to Claim 1, characterized in that the chiral auxiliary used is a (+)-L-tartaric acid derivative.
8. Process according to Claim 1, characterized in that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide), (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate or diethyl (+)-L-tartrate.
9. Process according to Claim 1, characterized in that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide) or (+)-L-tartaric acid bis-(N-morpholinamide).

**10.** Process according to Claim 1, characterized in that the oxidation is carried out in the presence of an organic base.

**11.** Process according to Claim 1, characterized in that the oxidation is carried out in the presence of a tertiary amine.

**12.** Process according to Claim 1, characterized in that the oxidation is carried out in organic solvents.

**13.** Process according to Claim 1, characterized in that the oxidation is carried out in organic solvents comprising 0 to 0.3% by volume of water.

**14.** Process according to Claim 1, characterized in that the oxidation is carried out in solvents of commercially available quality.

**15.** Process according to Claim 1, characterized in that the oxidation is carried out in an organic solvent which essentially comprises methyl isobutyl ketone.

**16.** Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide, or zirconium(IV) isopropoxide/isopropanol complex, that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide), (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate or diethyl (+)-L-tartrate.

**17.** Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide, or zirconium(IV) isopropoxide/isopropanol complex, that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide), (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate or diethyl (+)-L-tartrate, and that the oxidation is carried out in the presence of an organic base.

**18.** Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex, that the

chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide) or (+)-L-tartaric acid bis-(N-morpholinamide) and that the oxidation is carried out using cumene hydroperoxide.

19. Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex, that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide) or (+)-L-tartaric acid bis-(N-morpholinamide) and that the oxidation is carried out using cumene hydroperoxide in the presence of a tertiary amine.

20. (S)-pantoprazole prepared by the process according to Claim 1.

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
24 June 2004 (24.06.2004)

PCT

(10) International Publication Number  
**WO 2004/052881 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 401/12**

IX, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, US, VN, YU, ZA, ZW.

(21) International Application Number:

PCT/EP2003/013604

(22) International Filing Date: 3 December 2003 (03.12.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

02027274.6 6 December 2002 (06.12.2002) EP  
103 40 254.3 29 August 2003 (29.08.2003) DE

(71) Applicant (for all designated States except US): **ALTANA PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KOHL, Bernhard** [DE/DE]; Zum Brühl 9, 78465 Konstanz (DE). **MÜLLER, Bernd** [DE/DE]; Bücklestr. 84a, 78467 Konstanz (DE). **WEINGART, Ralf Steffen** [DE/DE]; Thingolstr. 34, 78465 Konstanz (DE).

(74) Agent: **WOLF, Ulrich**; Altana Pharma AG, Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(81) Designated States (national): AE, AL, AU, BA, BR, CA, CN, CO, DZ, EC, EG, GH, HR, ID, IL, IN, IS, JP, KR, LT,

(84) Designated States (regional): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

**Declarations under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AL, AU, BA, BR, CA, CN, CO, DZ, EC, EG, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, VN, YU, ZA, ZW; Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR)  
— of inventorship (Rule 4.17(iv)) for US only

**Published:**

— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

4 November 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PROCESS FOR PREPARING (S)-PANTOPRAZOLE**

(57) Abstract: The invention relates to a novel process for preparing (S)-pantoprazole using a chiral zirconium complex or a chiral hafnium complex.

**WO 2004/052881 A3**

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 03/13604

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category \* Citation of document, with indication, where appropriate, of the relevant passages

Relevant to claim No.

Y	WO 96/02535 A (COTTON HANNA KRISTINA ; LARSSON ERIK MAGNUS (SE); ASTRA AB (SE); SOERE) 1 February 1996 (1996-02-01) cited in the application examples 23,24	1-6, 10-15
X	example 23	20
Y	BONCHIO M ET AL: "The first Chiral Zirconium(IV) catalyst for highly stereoselective sulfoxidation" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY, EASTON, US, vol. 64, no. 4, 1999, pages 1326-1330, XP002242676 ISSN: 0022-3263 page 1327, scheme 2 page 1329, column 2, line 16 - page 1330, column 1, line 3	1-6, 10-15

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*Z\* document member of the same patent family

Date of the actual completion of the international search

9 September 2004

Date of mailing of the international search report

17/09/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx: 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hass, C



## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 03/13604

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92/08716 A (BYK GULDEN LOMBERG CHEM FAB) 29 May 1992 (1992-05-29) cited in the application examples 1-4 example 4 -----	1, 2
X	-----	20
X	WO 94/24867 A (SEPRACOR INC) 10 November 1994 (1994-11-10) cited in the application abstract page 2, line 7 - line 12 -----	20
A	WO 94/25028 A (SEPRACOR INC) 10 November 1994 (1994-11-10) cited in the application page 2, line 7 - line 12 -----	20
A	WO 99/47514 A (KNOLL AG ; BRENNAN JAMES PATRICK (GB); TURNER ANDREW TIMOTHY (GB)) 23 September 1999 (1999-09-23) examples 1,5 -----	1, 2
A	WO 96/17076 A (ASTRA AB ; HOLT ROBERT (GB); LINDBERG PER (SE); REEVE CHRISTOPHER (GB)) 6 June 1996 (1996-06-06) cited in the application page 5 page 7 page 13, line 15 - page 15, line 9 -----	1, 2

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 03/13604

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9602535	A	01-02-1996	SE 504459 C2
			17-02-1997
			AT 242233 T
			15-06-2003
			AU 688074 B2
			05-03-1998
			AU 2994895 A
			16-02-1996
			BR 9508292 A
			23-12-1997
			CA 2193994 A1
			01-02-1996
			CN 1157614 A , B
			20-08-1997
			CZ 9700064 A3
			11-06-1997
			DE 69530987 D1
			10-07-2003
			DE 69530987 T2
			19-05-2004
			DK 773940 T3
			15-09-2003
			EE 3354 B1
			15-02-2001
			EP 0773940 A1
			21-05-1997
			ES 2199998 T3
			01-03-2004
			FI 970102 A
			10-01-1997
			HK 1008331 A1
			21-11-2003
			HR 950401 A1
			31-10-1997
			HU 76642 A2
			28-10-1997
			IL 114477 A
			24-07-2001
			JP 10504290 T
			28-04-1998
			MA 23611 A1
			01-04-1996
			NO 970153 A
			14-01-1997
			NZ 289959 A
			26-01-1998
			PL 318165 A1
			26-05-1997
			PT 773940 T
			31-10-2003
			RU 2157806 C2
			20-10-2000
			SE 9402510 A
			16-01-1996
			WO 9602535 A1
			01-02-1996
			SI 773940 T1
			29-02-2004
			SK 4897 A3
			06-08-1997
			TR 960063 A2
			21-06-1996
			US 5948789 A
			07-09-1999
			ZA 9505724 A
			15-01-1996
WO 9208716	A	29-05-1992	DE 4035455 A1
			14-05-1992
			AU 8840691 A
			11-06-1992
			WO 9208716 A1
			29-05-1992
WO 9424867	A	10-11-1994	AU 6713194 A
			21-11-1994
			AU 6904498 A
			16-07-1998
			CA 2161256 A1
			10-11-1994
			EP 0695123 A1
			07-02-1996
			JP 8509736 T
			15-10-1996
			WO 9424867 A1
			10-11-1994
US			2003086968 A1
			08-05-2003
			US 2004142032 A1
			22-07-2004
			US 5888535 A
			30-03-1999
			US 2001008899 A1
			19-07-2001
WO 9425028	A	10-11-1994	AU 6713394 A
			21-11-1994
			CA 2161130 A1
			10-11-1994
			EP 0700292 A1
			13-03-1996
JP			8509738 T
			15-10-1996
			WO 9425028 A1
			10-11-1994
US			2002019420 A1
			14-02-2002
WO 9947514	A	23-09-1999	AU 3410699 A
			11-10-1999
			BR 9908835 A
			21-11-2000

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 03/13604

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9947514	A	CA 2323422 A1	23-09-1999
		CN 1293670 T	02-05-2001
		WO 9947514 A1	23-09-1999
		EP 1071678 A1	31-01-2001
		HU 0101230 A2	28-10-2001
		JP 2002506862 T	05-03-2002
		NO 20004580 A	14-09-2000
		SK 13452000 A3	09-04-2001
		TR 200002670 T2	21-11-2000
		TW 473476 B	21-01-2002
WO 9617076	A	06-06-1996	
		AT 232907 T	15-03-2003
		AU 699577 B2	10-12-1998
		AU 4126996 A	19-06-1996
		CA 2203999 A1	06-06-1996
		DE 69529686 D1	27-03-2003
		DE 69529686 T2	15-01-2004
		DK 795024 T3	22-04-2003
		EP 0795024 A1	17-09-1997
		ES 2191066 T3	01-09-2003
		JP 10510705 T	20-10-1998
		WO 9617076 A1	06-06-1996
		US 5840552 A	24-11-1998